

## THE EFFECTS OF ICE

often last for between 4 and 12 hours depending on how much is consumed.  
 usually felt quickly, however, it can take 1 to 2 days to entirely leave the body

### INTERMEDIATE

### LONG TERM

- Headaches and dizziness
- Dilated (enlarged) pupils and blurred vision
- Dry mouth
- Jaw clenching, teeth grinding
- Insomnia
- Increased attention, alertness and talkativeness

- Stroke
- Dental problems: increased tooth sensitivity, cracked teeth and cavities, gum disease
- Dependence
- Poor concentration and memory

- Trembling
- Sweaty, cold and clammy skin
- Increased breathing and body temperature
- Increased heart rate and blood pressure
- Reduced appetite
- Stomach cramps, nausea and vomiting
- Dehydration
- Aggressive behaviour
- Increased energy
- Changes in libido

- Increased risk of hepatitis C and HIV in injecting users
- Heart and lung problems, chest pains
- Kidney problems, including kidney failure
- Malnutrition and weight loss
- Exhaustion
- Movement problems



- A 'comedown' is often experienced when the drug wears off
- Symptoms can last for a few days and can include feeling down or depressed, exhausted and anxious.



- Withdrawals are unpleasant symptoms experienced by users who are dependent on ice and stop taking the drug
- Symptoms can last up to several weeks and can include headaches, cramps and vomiting, as well as anxiety, restlessness and aggression.



- Ice triggers the release of two chemicals in the brain, dopamine and noradrenaline
- These chemicals make us feel alert and excited
- This can have both short term and long term consequences
- For more information, see 'Factsheet: The effects of ice on the brain'

Last updated: Wednesday, 12 April 2018